

SBAR Tool

Postpartum Complications: Station 2

<p>Situation:</p>	<p>Jamie Miller, 40 y/o, rm 224; g 4 p 3 vaginal delivery 6 hours ago. HX: Shoulder dystocia</p> <p>Baby girl, apgar 6-8; 9lbs 2oz. Pediatrician Dr. Lim 41 wks gestation. Crepitates of right clavicle, no movement in left arm. Dr Lim notified, x-ray ordered.</p>
<p>Background:</p>	<p>Mother: Schedule induction at 41 wks. Bld type O + GBS: neg, Rubella: non-immune, Hepatitis neg, RPR: neg, HIV: neg.</p> <p>2 degree perineal laceration. Right labia hematoma. Pain level 6/10, medicated 30 min. ago.</p> <p>Plans on exclusively breastfeeding.</p> <p>Void X 1, 250 mL. VS. 118/78, 99.6, 101, 22</p> <p>Baby girl, 6 hour olds, voided at delivery, no stool, breast fed poorly, sleepy. VS 98.2, 158, 58 BS 45</p>
<p>Assessment:</p>	<p>Jamie is concerned about baby. She stated that all of her other deliveries were all so easy. She has never had so much difficulty or pain. She stated that nothing has gone the way she wanted it to. She needed Pitocin augmentation and then the baby's head got stuck. Everyone yelling at her, telling her to push harder.</p> <p>After delivery had perineal pain 10/10. Pain now 6/10. Jamie does not want to move.</p> <p>Fundus firm, flow moderate. QBL 700 mL.</p> <p>Baby girl, Haley, is sleepy, no interest in breastfeeding. Right arm secured. Right arm no movement, positive grasp.</p> <p>Father at bedside. Holding baby.</p>
<p>Recommendations:</p> <p><i>What are your concerns/recommendations?</i></p>	<p><u>Concerns:</u> <i>Mother: risk for PPH, PTSD and pain</i> <i>Baby: pain, feeding difficulty, hypoglycemia and assessment for brachial plexus of left arm.</i> <i>Delayed bonding.</i></p> <p><u>Recommendations:</u> <i>Mother: Ice, pain management, assessment of perineum, measuring QBL</i> <i>Baby: securing right arm, proper feeding position, monitoring pain, and glucose monitoring.</i></p>